



Pharmaceutical
Society of Australia



Sleeping Problems

Refreshing sleep is important for our emotional and physical wellbeing. Poor sleep can lead to drowsiness, poor concentration and other symptoms that interfere with daytime activities. The best way to treat sleeping problems is to find and manage the cause.

The amount of sleep we need depends on our age, lifestyle, diet, personality and circumstances. Most adults need 7 to 9 hours of sleep per night, teenagers need about 9.5 hours of sleep per night and infants need about 16 hours of sleep per day.

Sleep problems affect quality of sleep, amount of sleep or sleep behaviour. They include:

- Insomnia – insomnia is difficulty falling asleep or remaining asleep and/or not feeling refreshed by sleep. It is very common and may be a symptom of another condition
- Obstructive sleep apnoea – a condition causing a person to stop breathing for short periods while sleeping
- Restless legs syndrome – a condition causing an urge to move the legs when lying down
- Sleep behaviour disorders such as night terrors, sleep walking.


Signs of a sleep problem include:

- Finding it hard to go to sleep
- Restless sleep
- Waking up during the night, or very early in the morning, and not being able to get back to sleep
- Not feeling refreshed from sleep
- Feeling tired, irritable, anxious or depressed
- Finding it difficult to concentrate, remember things and make decisions.

What causes insomnia?

Things that can cause insomnia or make it worse include:

- Pain or discomfort
- Snoring (can be a sign of sleep apnoea)
- Stress, anxiety and emotional issues (e.g., unhappiness, fear, anger)
- Worrying about not sleeping well



- Having naps during the day
- Not enough physical exercise
- Sleeping in a room that is stuffy, noisy, or too brightly lit
- Sleeping in an uncomfortable bed or feeling too hot or too cold
- Going to bed with an empty or full stomach
- Being overstimulated in the evening by exercise, working, TV or radio
- Having caffeine (e.g., coffee, tea, chocolate, cola, energy drinks) in the evening
- Using nicotine (e.g., cigarettes, nicotine replacement therapy products)
- Using illicit drugs
- Drinking excess alcohol in the evening
- Changing work shifts or time zones (e.g., jet lag)
- Needing to pass urine often during the night (e.g., men with prostate disease)
- Pregnancy
- Some medicines (ask a pharmacist)
- Some medical conditions (e.g., anxiety, depression, other sleep problems, allergy).

Self care

Sleeping 'DOs':

- Remember that your sleep needs may change over time
- Sleep in a darkened, quiet room with plenty of fresh air
- Use a comfortable, supportive mattress and pillow
- Use clean bedding appropriate for the climate
- Learn and practice relaxation techniques to manage stress
- Get enough exercise during the day to feel physically tired each night. (Avoid strenuous exercise within 3 hours of bedtime.)
- Limit daily caffeine consumption
- Limit daily alcohol consumption
- Go to bed at the same time each night and get up at the same time each day (trains the body to develop a regular sleep cycle)
- Have a calming bedtime routine such as:
 - A warm bath
 - A warm milk drink or a light snack
 - Listening to peaceful music and/or reading
 - Slow, deep breathing.

Sleeping 'DON'Ts':

- Don't nap during the day, especially late in the day
- Don't work or watch TV in your bedroom. Keep your bedroom for sleeping, relaxing and intimacy
- Don't have pets or highly illuminated equipment in the bedroom
- Don't have a heavy meal within 3 hours of bedtime
- Don't drink a lot of liquid in the evening (to help limit night time visits to the toilet)
- Don't have caffeine close to bedtime
- Don't drink alcohol close to bedtime (alcohol can affect sleep quality)
- Don't smoke.

Getting to sleep

If you cannot get to sleep, or if you wake up and cannot get back to sleep, get out of bed and:

- Write down worries or problems and decide to deal with them in the morning
- Do something relaxing or perhaps repeat your calming bedtime routine
- Return to bed when feeling sleepy.

Your **Self Care** Pharmacist

Important

Consult a doctor or pharmacist if you:

- Can't sleep due to illness, pain or discomfort
- Wake up coughing, wheezing or unable to breathe
- Snore, especially if you wake yourself or others
- Have been taking sleeping tablets regularly for more than four weeks
- Continue to be tired during the day.

Medicines

The best way to treat sleeping problems is to find and manage the cause. Sometimes sleeping tablets are also needed to help correct a sleep problem. Most sleeping tablets should be used at the lowest possible dose for the shortest possible time (preferably less than two weeks). Ask a doctor or pharmacist for advice.

- It is possible to become dependent on some sleeping tablets if they are used continuously.
- It is possible to become tolerant to some sleeping tablets, so that you need to take more and more to get the same effect.
- Sleeping tablets can cause daytime drowsiness, confusion and unsteadiness, especially in older people.
- Sleeping tablets may increase the effects of alcohol. The combination of sleeping tablets and alcohol should be avoided.
- Some herbal and complementary medicines may help relaxation and sleep.

Related fact cards

- *Alcohol*
- *Anxiety*
- *Bladder and Urine Control*
- *Depression*
- *Pain Relievers*
- *Post-Traumatic Stress Disorder*
- *Prostate Problems*
- *Relaxation Techniques*
- *Smoking series*

For more information

A doctor – listed under Medical Practitioners in the yellow pages of the phone book.

Australasian Sleep Association – www.sleep.org.au

HEALTHInsite – website www.healthinsite.gov.au

Consumer Medicine Information (CMI) leaflets – your pharmacist can advise on availability.

NPS Medicines Line – phone 1300 888 763 Monday to Friday, 9am to 6pm EST.

The Poisons Information Centre – in case of poisoning phone 131 126 from anywhere in Australia.

Pharmacists are medicines experts. Ask a pharmacist for advice when choosing a medicine.

Sponsored by



Australian Government

Department of Veterans' Affairs